

Brand name: ZENOL Pain Relief Patch

**RISK CATEGORIES**  
Third-class OTC drug

**CHARACTERISTICS**

- Topical analgesic anti-inflammatory agent

1. "The patch exerts high anti-inflammatory and cooling effects."  
With anti-inflammatory effect of glycolsalicylate and cooling effect of high moisture base, the patch is very effective for joint pain, muscle pain, and sprain.
2. "It stretches well and adhesive to the affected area."  
Using highly stretchable nonwoven fabric and highly viscous plaster, the patch exerts its analgesic effect by adhering to areas such as elbow, knee, etc. with a wide range of motion. "Aluminum bag with a zipper" The aluminum bag with a zipper helps easy storage, and prevents the plaster from being dried.



**ACTIVE INGREDIENTS**  
Per 100 g plaster (per 875 m2)

For further information:

		Distributed by	TAIHO PHARMACEUTICAL CO., LTD.	
<b>ACTIVE INGREDIENTS</b>	Content	Manufactured and Distributed by	Mikasa Seiyaku Co., Ltd.	
Glycol Salicylate	1.70 g	Dosage form	Pap	
L-Menthol	0.50 g	Packaging unit	1 box (14 cm x 10 cm x 24 sheets)	1 box (14 cm x 10 cm x 24 sheets) x 2 boxpackage
Tocopherol Acetate	0.30 g	Manufacturer's suggested retail price	JPY 2,200 (JPY 2,000 excluding tax)	JPY 4,400 (JPY 4,000 excluding tax)
Inactive ingredients: Gelatin, Propylene Glycolate, Isopropyl Myristate, Polysorbate 80, Disodium Edetate, Paraben, Glycerin, Partially Neutralized Polyacrylate, Polyacrylate, Carmellose Sodium, Tartaric Acid, Caster Oil, Propylene Glycol Fatty Acid Ester, Acrylic Acid Starch, Titanium Oxide, Aluminum Hydroxide		JAN Code	4987117372151	4987117372182
		Expiration	3 years	

**DOSAGE AND ADMINISTRATION**  
Remove the film on the surface, and apply an appropriate amount to the affected area once to twice daily.

<Precautions>

- (1) Strictly follow the dosing instructions.
- (2) For children, use under parental guidance.
- (3) Wipe off sweat on the affected area before use.

## INDICATIONS

Joint pain, muscle pain, sprain, stiff shoulder, muscle fatigue, low back pain, bruise, fracture pain, frostbite

## PRECAUTIONS FOR USE

☒ The following precaution should be observed.

(If not observed, it may aggravate the current symptoms or increase the risk of adverse reactions.)

Do not apply to the following areas.

- (1) Around the eyes, mucous membranes, etc.
- (2) Eczema, contact dermatitis, wounds



### Consultation

1. The following persons should consult a physician, pharmacist or registered salesperson before using the patch. Persons with a history of allergic symptoms due to any drug
2. The following symptoms may be adverse reactions to the patch. If any of these symptoms occur, discontinue the use of the patch immediately, and consult a physician, pharmacist or registered salesperson with this leaflet.

Affected body system	Symptoms
Skin	Rash, redness, itching

3. If symptoms do not improve after using the patch for five to six days, discontinue the use of the patch, and consult a physician, pharmacist or registered salesperson with this leaflet.

## CONTACT

For inquiries, please contact the shop where you purchased the product or the following:

Distributor Contact: Customer Relations Office, Taiho Pharmaceutical Co., Ltd.

1-27 Kandnishiki-cho, Chiyoda-ku, Tokyo, 101-8444, Japan

Phone: 0120-4527-66

Business hours: 9:00-17:00 (Excluding weekends and holidays)

Website: <https://www.taiho.co.jp/>

Manufacturer and distributor : MIKASA SEIYAKU CO., LTD