Brand name: ZENOL XUM SX

RISK CATEGORIES

Second-class OTC drug

CHARACTERISTICS

Topical analgesic anti-inflammatory agent Felbinac penetrates into the affected area, and helps relieve joint, muscle, and shoulder pains. This is a stick which is easily applied to the affected area without staining hands. As it is not sticky, it allows putting on clothes

Easy-to-use new designed container





ACTIVE INGREDIENTS

immediately after use.

Per 100 g

	For	further	informa	ation:
--	-----	---------	---------	--------

		Distributed by:	TAIHO PHARMACEUTICAL CO., LTD.
ACTIVE INGREDIENTS	Content	Manufactured and Distributed by:	Mikasa Seiyaku Co., Ltd.
Felbinac 3.0 g		Dosage form	Stick type agent
L-Menthol	3.0 g	Packaging unit	43 g x 1 box
Inactive ingredients: Isopropanol, Propylene Glycol, Macrogol, Polyoxyethylene Hardened		Manufacturer's suggested retail price	JPY 1,650 (JPY 1,500 excluding tax)
Caster Oil, Sodium St Oleyl Alcohol, Squalar	earate, 1,3-Butylene Glycol, ne, pH Adjuster	JAN Code	4987117372014
		Expiration	3 years

DOSAGE AND ADMINISTRATION

Apply an appropriate amount to the affected area twice to four times daily.

<Precautions>

- (1) Strictly follow the dosing instructions.
- (2) Be careful not to get the agent into the eyes. If the agent gets into the eyes, immediately wash with water or lukewarm water. If symptoms are severe, consult an ophthalmologist.
- (3) For external use only. Do not take the agent orally.
- (4) Do not cover the affected area with a poorlyventilatedmaterial such as wrap film, etc. after applying the agent.
- (5) Wipe off sweat on the affected area before use.
- (6) Twist the container until 4 to 5 mm of agent appears before use, to prevent the container from directly touching the affected area.
- (7) Stop using the agent when the bottom of the container appears and the upper side of the bottom is exposed. Do not forcibly use the agent remaining in the bottom of the container.

INDICATIONS

Joint pain, muscle pain, shoulder pain du shoulder, low back pain, bruise, sprain, tenosynovitis (pain and swelling in hands and ankles), elbow pain (tennis elbow, et

PRECAUTIONS FOR USE

▼ The following precaution should be observed.

(If not observed, it may aggravate the current symptoms or increase the risk of adverse reactions.)

- 1. The following persons should not use the agent.
- (1) Persons with a history of allergic symptoms to the agent or any of the ingredients of the agent
- (2) Persons with a history of asthma
- (3) Pregnant women or possibly pregnant women
- (4) Children under 15 years
- 2. Do not apply to the following areas.
- (1) Around the eyes, mucous membranes, etc.
- (2) Eczema, contact dermatitis, wounds
- (3) Athlete's foot (tinea pedis), tinea cruris, etc. or purulent areas

Consultation

- 1. The following persons should consult a physician, pharmacist or registered salesperson before using the agent.
- (1) Persons under treatment by a physician
- (2) Persons with a history of allergic symptoms to any drug
- 2. The following symptoms may be adverse reactions to the agent. If any of these symptoms occur, discontinue the use of the agent immediately, and consult a physician, pharmacist or registered salesperson with this leaflet.

Affected body system	Symptoms
Skin	Rash, redness, itching, swelling, tingling sensation, contact dermatitis

The following serious symptom may rarely occur. If the symptom occurs, consult a physician immediately.

Symptom	Symptoms
Shock (anaphylaxis)	Skin itching, urticaria, hoarse voice, sneezing, larynx itching, difficulty breathing, palpitation, consciousness clouding, etc. occur immediately after using the agent.

3. If symptoms do not improve after using the agent for about one week, discontinue the use of the agent, and consult a physician, pharmacist or registered salesperson with this leaflet.

CONTACT

For inquiries, please contact the shop where you purchased the product or the following:

Distributor Contact: Customer Relations Office, Taiho Pharmaceutical Co., Ltd.

1-27 Kandanishiki-cho, Chiyoda-ku, Tokyo, 101-8444, Japan

Phone: 0120-4527-66

Business hours: 9:00-17:00 (Excluding weekends and holidays)

Website: https://www.taiho.co.jp/

Manufacturer and distributor: MIKASA SEIYAKU CO., LTD