



Brand name: Zenol Tick E			
RISK CATEGORIES Third-class OTC drug CHARACTERISTICS <ul style="list-style-type: none"> ● Topical analgesic anti-inflammatory agent Methyl salicylate is effective for stiff shoulder and muscle fatigue by penetrating into the affected areas. This is a stick which is easily applied to the affected area without staining hands. As it is not sticky, it allows putting on clothes immediately after use. It has a cool flavor.			
ACTIVE INGREDIENTS <ul style="list-style-type: none"> ● Per 100 g 		For further information:	
		Distributed by:	TAIHO PHARMACEUTICAL CO., LTD.
ACTIVE INGREDIENTS	Content	Manufactured and Distributed by:	Mikasa Seiyaku Co., Ltd.
Methyl Salicylate	17.50 g	Dosage form	Stick type agent
L-Menthol	6.00 g	Packaging unit	33 g x 1 box
dl-Camphor	5.00 g	Manufacturer's suggested retail price	JPY 935 (JPY 850 excluding tax)
Glycyrrhetic Acid	0.02 g	JAN Code	4987117372052
Inactive ingredients: Isopropanol Flavor, Sodium Stearate, Glycerin, Caster Oil		Expiration	3 years
INDICATIONS Stiff shoulder, muscle fatigue, muscle pain, low back pain, joint pain, bruise, sprain, fracture pain, frostbite		DOSAGE AND ADMINISTRATION Usually, apply an appropriate amount to the affected area once to several times daily. <Precautions> (1) Strictly follow the dosing instructions. (2) Be careful not to get the agent into the eyes. If the agent gets into the eyes, immediately wash with water or lukewarm water. If symptoms are severe, consult an ophthalmologist. (3) For children, use under the supervision of their guardians. (4) For external use only. Do not take the agent orally. (5) Wipe off sweat on the affected area before use. (6) Twist the container until 4 to 5 mm of agent appears before use, to prevent the container from directly touching the affected area. (7) Stop using the agent if the bottom of the container appears and becomes unstable. Do not forcibly use the agent remaining in the bottom of the container.	
PRECAUTIONS FOR USE ☒ The following precaution should be observed. (If not observed, it may aggravate the current symptoms or increase the risk of adverse reactions.) Do not apply to the following areas. (1) Around the eyes, mucous membranes, etc.			

(2) Eczema, contact dermatitis, wounds

 Consultation

1. The following persons should consult a physician, pharmacist or registered salesperson before using the agent.
Persons with a history of allergic symptoms due to any drug
2. The following symptoms may be adverse reactions to the agent. If any of these symptoms occur, discontinue the use of the agent immediately, and consult a physician, pharmacist or registered salesperson with this leaflet.

Affected body system	Symptoms
Skin	Rash, redness, itching

3. If symptoms do not improve after using the agent for five to six days, discontinue the use of the agent, and consult a physician, pharmacist or registered salesperson with this leaflet.