

Brand name: Taiho Oral Solution for Cold “Gingyo-san”

## RISK CATEGORIES

Second-class OTC drug

## CHARACTERISTICS

Taiho Oral Solution for Cold “Gingyo-san” is an oral solution for cold prepared based on “Gingyo-san” listed in a Chinese medical book, “the Wen Bing Tiao Bian (Systematic Differentiation of Febrile Diseases)”. “Gingyo-san” containing 10 types of crude drugs including lonicerae flos, forsythiae fructus, saigae tataricae cornu, etc. is effective for the treatment of common cold symptoms including throat pain, cough, headache, etc.



## ACTIVE INGREDIENTS

Per 90 mL (30 mL x 3 bottles) daily

Gingyo-san Extract 9.7 g

As crude drugs,

Lonicerae Flos ..... 4.260 g forsythiae

Fructus ..... 4.260 g

Platycodon Root..... 2.556 g

Glycyrrhiza ..... 2.556 g

Mentha Herb ..... 2.556 g

Fermented Soybeans ..... 2.136 g

Burdock Fruit ..... 2.136 g

Lophatheri Herba ..... 1.704 g

Schizonepeta Spike ..... 1.704 g

Saigae Tataricae Cornu ..... 0.132 g

Inactive ingredients: White Soft Sugar, High-fructose Corn Syrup, Xylitol, Citric Acid, Sodium Citrate, Sodium Benzoate, Paraben, Polyoxyethylene Hardened Caster Oil,, Alcohol, Flavor (l-Menthol)

### <Precautions>

Shake well before use. As this medicine contains crude drugs, it may cause precipitation.

For further information:

Manufactured and Distributed by:

TAIHO PHARMACEUTICAL CO., LTD.

Dosage form

Liquid

Packaging unit

30 mL x 1 bottle

30 mL x 3 bottles

Manufacturer's suggested retail price

JPY 341  
(JPY 310 excluding tax)

JPY 1,023  
(JPY 930 excluding tax)

JAN Code

45144677

4987117234619

Expiration

3 years

## DOSAGE AND ADMINISTRATION

Take the following dose between meals after shaking well.  
(Between meals mean 2 to 3 hours after meals. )

Age

Adults (15 years and over):

Children (under 15 years)

Dose

1 bottle

Daily dose

3 times  
(between meals)

Do not use.

## INDICATIONS

Throat pain, cough, dry mouth (throat), and headache due to common cold

### <Precautions>

(1) Strictly follow the dosing instructions.

(2) Take one full bottle as a single dose.

## PRECAUTIONS FOR USE

☒ The following precaution should be observed.

(If not observed, it may aggravate the current symptoms or increase the risk of adverse reactions.)

The duration of treatment should be limited to short term, and continuous treatment should be avoided.

### 📖 Consultation

1. The following persons should consult a physician, pharmacist or registered salesperson before using the medicine.

- (1) Persons under treatment by a physician
- (2) Pregnant women or possibly pregnant women
- (3) Persons who are physically weak (declining of strength or constitutional weakness)
- (4) Persons with gastrointestinal weakness
- (5) Elderly
- (6) Persons with a history of rash, redness, itching, etc. due to any drug
- (7) Persons with the following symptom: oedema
- (8) Persons diagnosed with the following diseases: hypertension, cardiac disease or kidney disease

2. The following symptoms may be adverse reactions to the medicine. If any of these symptoms occur, discontinue the use of the medicine immediately, and consult a physician, pharmacist or registered salesperson with this medicine.

Affected body system	Symptoms
Skin	Rash, redness, itching
Gastrointestinal	Feeling queasy, loss of appetite, stomach discomfort

The following serious symptoms may rarely occur. If the symptom occurs, consult a physician immediately.

Symptom	Symptoms
Pseudoaldosteronism, myopathy	Weakness, numbness, tightness, and stiffness of limbs occur along with feelings of weakness and muscle pain, and they gradually get worse.

3. If symptoms do not improve after using the medicine for five to six times, discontinue the use of the medicine, and consult a physician, pharmacist or registered salesperson with this medicine.