

Brand name: Harnicare Bearberry Tablets

**RISK CATEGORIES**  
Second-class OTC drug

**CHARACTERISTICS**  
Harnicare-series tablet focusing on easing inflammation of the urinary bladder.  
Bearberry has been tableted for the first time in Japan.  
The active ingredient, bearberry, is contained in the maximum amount in this OTC drug.  
The pink-based package easily attracts the attention of women.



**ACTIVE INGREDIENTS** (in a daily dose of 12 tablets)  
Dried bearberry extract: 1,875 mg (equivalent to 15 g bearberry)

[Inactive ingredients]  
Dextrin, crystalline cellulose, potato starch, croscarmellose sodium, light anhydrous silicic acid, magnesium stearate

For further information:

Distributed by:	TAIHO PHARMACEUTICAL CO., LTD.		
Dosage form	Tablet		
Packaging unit	48 tablets	84 tablets	
Manufacturer's suggested retail price	2,024 JPY (Price before tax: 1,840 JPY)	3,365 JPY (Price before tax: 3,059JPY)	
JAN code	4987117613100	4987117613209	
Expiration	3 years		

**DOUSAGE AND ADMINISTRATION**  
Take the dose indicated below with cold or warm water before a meal or between meals.

Age	Single dose	Daily dose
Adult (15 years of age or older)	4 tablets	3 times
Under 15 years of age	Do not take this product.	

**INDICATIONS**  
Feeling of incomplete emptying (difficulty in urination)  
Discomfort during urination

Note: 'Between meals' means about 2 hours after a meal.  
● Strictly follow the prescribed dosage and administration.

**PRECAUTIONS FOR USE AND CONSULTATIONS**

- Those described below should consult with a doctor, pharmacist, or registered distributor before use:
  - Those who are currently seeking medical treatment;
  - Women who are or may be pregnant;
  - Those with a sensitive stomach;
  - Those who have developed allergy symptoms due to drugs or the like; or
  - Those who are diagnosed with the following disease: Renal dysfunction.
- Developing any of the symptoms listed below after taking this product may indicate the occurrence of an adverse reaction. If so, immediately stop taking the product and consult with a doctor, pharmacist, or registered distributor, showing the product package.

Related organ	Symptom
Skin	Rash, redness, itching
Digestive organs	Stomach discomfort, nausea and vomiting

- If your symptom does not resolve after one month, stop taking the product and consult with a doctor, pharmacist, or registered distributor, showing the product package.
- If you wish to continue taking this product for an extended period of time, consult with a doctor, pharmacist, or registered distributor.

**CONTACT**

For inquiries, please contact the shop where you purchased the product or the following:

Manufacturer:

Customer Relations Office, Taiho Pharmaceutical Co., Ltd.

1-27 Kandanshiki-cho, Chiyoda-ku, Tokyo, 101-8444, Japan

Phone: 0120-4527-66 Business hours: 9:00-17:00 (Excluding weekends and holidays)

Website: <https://www.taiho.co.jp/>