


Brand name: Harncare Drink																					
<p>RISK CATEGORIES Designated second-class OTC drug</p> <p>CHARACTERISTICS</p> <ul style="list-style-type: none"> This is a medicine prepared using an original formulation method and specialized for the treatment of urinary symptoms. <p>(1) This is an herbal medicine containing extracts prepared by extracting and concentrating eight crude drugs (Rehmannia Root, Alisma Tuber, Moutan Bark, Poria Sclerotium, Cornus Fruit, Dioscorea Rhizome, Cinnamon Bark, and Roasted Aconite), adding ethanol, separating and removing starch, and evaporating and removing ethanol.</p> <p>(2) The medicine relieves mild incontinence, pollakiuria (frequent urination), feeling of residual urine, and vesical tenesmus accompanied by decreased physical strength, decline of the lower body or cold limbs.</p> <p>(3) This is easy-to-take liquid.</p>																					
	<p>ACTIVE INGREDIENTS</p> <p>Per 2 bottles (60 mL) daily: Crude drug extract H.....11 mL Extracted from Rehmannia Root...5 mg, Poria Sclerotium...3 g, Cinnamon Bark...1 g, Alisma Tuber...3 g, Cornus Fruit... 3 g, Roasted Aconite... 1 g, Moutan Bark...3 g, and Dioscorea Rhizome...3 g</p> <p>Inactive ingredients: Glycerin, Polyoxyethylene Hardened Caster Oil, Citric Acid, Sodium Benzoate, Paraben, pH-Adjuster, Corn Starch, Flavor .. Shake well before use. This Patch may cause precipitation because it contains crud drug components.</p>		<p>For further information:</p> <table border="1"> <tr> <td>Manufactured and Distributed by:</td> <td colspan="2">TAIHO PHARMACEUTICAL CO., LTD.</td> </tr> <tr> <td>Dosage form</td> <td colspan="2">Liquid</td> </tr> <tr> <td>Packaging unit</td> <td>30 mL x 2 bottles</td> <td>30 mL x 10 bottles</td> </tr> <tr> <td>Manufacturer's suggested retail price</td> <td>JPY 968 (JPY 880 excluding tax)</td> <td>JPY 4,598 (JPY 4,180 excluding tax)</td> </tr> <tr> <td>JAN Code</td> <td>4987117612028</td> <td>4987117612219</td> </tr> <tr> <td>Expiration</td> <td colspan="2">3 years</td> </tr> </table>		Manufactured and Distributed by:	TAIHO PHARMACEUTICAL CO., LTD.		Dosage form	Liquid		Packaging unit	30 mL x 2 bottles	30 mL x 10 bottles	Manufacturer's suggested retail price	JPY 968 (JPY 880 excluding tax)	JPY 4,598 (JPY 4,180 excluding tax)	JAN Code	4987117612028	4987117612219	Expiration	3 years
Manufactured and Distributed by:	TAIHO PHARMACEUTICAL CO., LTD.																				
Dosage form	Liquid																				
Packaging unit	30 mL x 2 bottles	30 mL x 10 bottles																			
Manufacturer's suggested retail price	JPY 968 (JPY 880 excluding tax)	JPY 4,598 (JPY 4,180 excluding tax)																			
JAN Code	4987117612028	4987117612219																			
Expiration	3 years																				
<p>INDICATIONS</p> <p>It relieves the following symptoms accompanied by decreased physical strength, decline of the lower body or cold limbs. Mild incontinence, pollakiuria (frequent urination), feeling of residual urine, vesical tenesmus</p>	<p>DOSAGE AND ADMINISTRATION</p> <p>Take the following dose before breakfast and dinner or between meals.</p> <table border="1"> <tr> <td>Age</td> <td>Adults (15 years and over)</td> <td>Children (under 15 years)</td> </tr> <tr> <td>Dose</td> <td>1 bottle (30 mL)</td> <td rowspan="2">Do not use.</td> </tr> <tr> <td>Daily dose</td> <td>Twice daily</td> </tr> </table>		Age	Adults (15 years and over)	Children (under 15 years)	Dose	1 bottle (30 mL)	Do not use.	Daily dose	Twice daily											
	Age	Adults (15 years and over)	Children (under 15 years)																		
Dose	1 bottle (30 mL)	Do not use.																			
Daily dose	Twice daily																				
<p>Note:.) Between meals mean about 2 hours after meals. ● Strictly follow the prescribed dosing instructions.</p>																					

PRECAUTIONS FOR USE

☒ The following precaution should be observed.

(If not observed, it may aggravate the current symptoms or increase the risk of adverse reactions.)

The following persons should not take this medicine.

(1) Persons with gastrointestinal weakness

(2) Persons who are prone to diarrhea

(3) Persons with the following symptoms

- “Not being aware of incontinence” due to spinal cord injury, dementia, etc.
- “Always having a small amount of incontinence” due to prostate hyperplasia, etc.

Consultation

1. The following persons should consult a physician, pharmacist or registered salesperson before using this medicine.

(1) Persons under treatment by a physician

(2) Pregnant or possibly pregnant women

(3) Persons with strong constitution, severe hot flush, and red face

(4) Persons with a history of rash, redness, itching, etc. due to any drug

(5) Persons using other Kampo preparations (Caution should be exercised to duplicate use of crude drug-containing products.)

2. The following symptoms may be adverse reactions to the medicine. If any of these symptoms occur, discontinue the use of the medicine immediately, and consult a physician, pharmacist or registered salesperson with this leaflet.

Affected body system	Symptoms
Skin	Rash, redness, itching
Gastrointestinal	Feeling queasy/vomiting, loss of appetite, stomach discomfort, diarrhea, abdominal pain, constipation
Psychoneurotic	Headache, dizziness
Cardiovascular	Palpitation
Respiratory	Shortness of breath
Urinary	Urinary retention
Others	Hot flush, chills, oedema, numbness of lip/tongue

3. If symptoms do not improve after using the medicine for about two weeks, discontinue the use of this medicine, and consult a physician, pharmacist or registered salesperson with this leaflet.