

Brand Name: Pitas Cough Troche

**RISK CATEGORIES**

Second-class OTC drugs

**CHARACTERISTICS**

- A cough suppressant ingredient, dextromethorphan phenolphthalinate, acts on the cough center and controls painful coughs
- An expectorant, potassium guaiacolsulfonate, suppresses phlegm
- Can be taken without water
- A slim package makes it convenient to carry around



**ACTIVE INGREDIENTS (In 6 troches)**

Dextromethorphan Phenolphthalinate 60 mg  
Potassium Guaiacolsulfonate 140 mg  
Cetylpyridinium Chloride Hydrate 6 mg

[Inactive ingredients]: Hydroxypropylcellulose, Povidone, Pullulan, Macrogol, l-Menthol, Tannic Acid, D-Sorbitol, Sodium Saccharine, Sucrose Fatty Acid Ester, Sucralose, Sodium Copper Chlorophyllin, Flavor

For further information:

Distributed by	TAIHO PHARMACEUTICAL CO., LTD.
Manufactured by	KYUKYU PHARMACEUTICAL CO., LTD.
Dosage form	Troche
Packaging unit	12 troches
Manufacturer's suggested retail price	JPY 770 (JPY 700 excluding tax)
JAN Code	45198625
Expiration date	3 years

**DOSAGE AND ADMINISTRATION**

Put the following dose in the mouth, and slowly dissolve without chewing.

Dose for Adults (15 years and over): 1 troche

Frequency of daily dosing: 6 times

Dosing interval: 2 hours or longer

Do not use for children under 15 years of age

**<Precautions>**

(1) Strictly follow the dosing instructions.

(2) Do not chew or swallow whole.

**INDICATIONS**

Cough

Phlegm

Throat pain, throat swelling, sore throat, throat discomfort, and hoarse voice due to throat inflammation

**PRECAUTIONS FOR USE**

The precaution should be observed.

(If not observed, it may aggravate the current symptoms or increase the risk of adverse reactions.)

1. The following persons should not take this medicine:

- Persons having a history of allergy to this product or any component of this product.

2. During use of this product, the following drugs may not be used.

Other antitussive expectorants, cold medicines, sedatives, oral-dose drugs containing antihistamine, and so on (oral-dose drugs for rhinitis, drugs for motion sickness, anti-allergy drugs, etc.)

**Consultation**

1. The following persons should consult a physician, pharmacist or registered salesperson before using this medicine.

- (1) Persons under treatment by a physician

- (2) Pregnant or possibly pregnant women
- (3) Persons having a history of allergy to drugs, etc.
- (4) Persons having any of the following symptoms: High fever

2. The following symptoms may be adverse reactions to the medicine. If any of these symptoms occur, discontinue the use of this medicine immediately, and consult a physician, pharmacist or registered salesperson with this leaflet.

Affected area	Skin	Gastrointestinal System	Neuropsychiatric System	Respiratory System
Symptoms	Eruption/redness, itching sensation	nausea/vomiting, anorexia	dizziness	dyspnea, shortness of breath

Although rarely, the following serious symptoms may appear. In such an event, immediately consult a physician.

Symptom name	Features
Shock (anaphylaxis)	Immediately after intake of the product, there may appear an itching sensation, urticaria, hoarse voice, sneezing, throat pruritus, dyspnea, palpitation, consciousness clouding, etc

3. If the symptoms do not improve after using the medicine for 5-6 times, discontinue the use of this medicine, and consult a physician, pharmacist or registered salesperson with this leaflet.