Brand name: PITAS Medical Troche

## RISK CATEGORIES

Designated quasi-drug

## CHARACTERISTICS

- Handy product for sterilization of the mouth that can be taken without water
- Plum flavor
- -A slim package fits in a card case to carry around.





#### ACTIVE INGREDIENTS (in 6 pieces)

Cetylpyridinium Chloride Hydrate: 8.28 mg

[Inactive ingredients] Hydroxypropylcellulose, Sucralose, Povidone, Pullulan, Macrogol, I-Menthol, Tannic Acid, D-Sorbitol, Saccharin Sodium, Sucrose Fatty Acid Ester, Citric Acid Hydrate, New Coccine, Ethyl Vanillin, Vanillin, Flavor

#### For further information:

Manufactured and distributed by:	TAIHO PHARMACEUTICAL CO., LTD.		
Dosage form	Troche		
Packaging unit	12 troches		
Manufacturer's suggested retail price	JPY 696 (Price before tax: JPY 633)		
JAN code	4987117220025		_
Expiration	3 years		

## DOSAGE AND ADMINISTRATION

Put the following dose in the mouth, and slowly dissolve without chewing.

Dose for Adults (15 years and over): 1 troche for a single dose Frequency of daily dosing: 4 to 6 times
Do not use for children under 15 years old

# **■**INDICATIONS

- Sterilization and disinfection of the mouth
- Throat pain, throat swelling, sore throat, throat discomfort, and hoarse voice due to throat inflammation
- Removal of bad breath

- <Precautions for the dosage and administration>
- (1) Strictly follow the dosage and administration instructions.
- (2) Do not chew or swallow it.

# PRECAUTIONS FOR USE AND CONSULTATIONS

- 1. Those described below should consult with a doctor, dentist, or pharmacist before use.
  - (1) Those who are currently seeking medical or dental treatment;
  - (2) Women who are or may be pregnant;
  - (3) Those who have an allergic reaction or whose family member has an allergic reaction; or
  - (4) Those who have developed allergy symptoms due to drugs or the like.
- 2. In the cases described below, immediately stop using the product and consult with a doctor, dentist,
- or pharmacist, showing these instructions.
  - (1) You develop any of the symptoms listed below after using this product: Skin: Rash, redness, itching
  - (2) Your symptom does not improve after one week.

## PRECAUTIONS FOR STORAGE AND HANDLING

- (1) Store in a cool, dry place, away from direct sunlight.
- (2) Keep out of reach of children.
- (3) Do not transfer the product to a different container. (It may lead to misuse or alter the quality of the product.)
- (4) Use immediately after opening an aluminum bag.
- (5) Do not use the product past the expiration date.

## CONTACT

For inquiries, please contact the shop where you purchased the product or the following:

Manufacturer and distributor

Contact: Customer Relations Office, Taiho Pharmaceutical Co., Ltd. 1-27 Kanda Nishiki-cho, Chiyoda-ku, Tokyo, 101-8444, Japan

Phone: 0120-4527-66

Business hours: 9:00 to 17:00 (Excluding weekends and holidays)

Website: <a href="https://www.taiho.co.jp/">https://www.taiho.co.jp/</a>