

Brand name: PITAS Throat Troche O (Orange Flavor)

RISK CATEGORIES
Designated quasi-drug

CHARACTERISTICS

A troche which sticks in the mouth to relieve throat pain and swelling. Able to intake while speaking.
 - Able to intake without water even in the middle of a meeting.
 - No active ingredients to cause drowsiness. Able to intake at work.
 - A slim package fits in a card case to carry around.



ACTIVE INGREDIENTS (in 6 troches)
Cetylpyridinium Chloride Hydrate 8.28mg
 [Inactive ingredients] Hydroxypropylcellulose, Sucralose, Povidone, Pullulan, Macrogol, l-Menthol, Tannic Acid, D-Sorbitol, Saccharin Sodium, Sucrose Fatty Acid Ester, dl- α -tocopherol, Flavor, New Coccine, Sunset Yellow FCF

For further information:

Manufactured and Distributed by:	TAIHO PHARMACEUTICAL CO., LTD.		
Dosage form	Troche		
Packaging unit	12 troches		
Manufacturer's suggested retail price	JPY 633 (JPY 575 excluding tax)		
JAN Code	45209604		
Expiration	3 years		

DOSAGE AND ADMINISTRATION

Put the following dose in the mouth, and slowly dissolve without chewing.
 Dose for Adults (15 years and over) 1 troche
 Frequency of daily dosing 4 to 6 times
 Do not use for children under 15

<Precautions>

- (1) Strictly follow the dosing instructions.
- (2) Do not chew or swallow whole.

INDICATIONS

Throat pain, throat swelling, sore throat, throat discomfort, and hoarse voice due to throat inflammation, sterilization/disinfection of the mouth, and removal of bad breath

PRECAUTIONS FOR USE

Consultation

1. The following persons should consult a physician, dentist or pharmacist before using the medicine.

- (1) Persons under treatment by a physician or dentist.
- (2) Pregnant women or possibly pregnant women.
- (3) The person who will take this drug or any on in the family has allergic constitution.
- (4) Persons with a history of allergic symptoms to any drug.

2. If any of these symptoms occur, discontinue the use of the medicine immediately, and consult a physician, dentist or pharmacist with this leaflet.

(1) If any of the following symptoms occur.

Skin: Rash, redness, itching

(2) If symptoms do not improve after using the medicine for one week.

CONTACT

For inquiries, please contact the shop where you purchased the product or the following:

Distributor

Contact: Customer Relations Office, Taiho Pharmaceutical Co., Ltd.

1-27 Kandanishiki-cho, Chiyoda-ku, Tokyo, 101-8444, Japan

Phone: 0120-4527-66

Business hours: 9:00-17:00 (Excluding weekends and holidays)

Website: <https://www.taiho.co.jp/>