


Brand name: Pitas Throat Troche																									
<p>RISK CATEGORIES Designated quasi-drug</p> <p>CHARACTERISTICS A troche which sticks in the mouth to relieve throat pain and swelling. Able to intake while speaking. - Able to intake without water even in the middle of a meeting. - No active ingredients to cause drowsiness. Able to intake at work. - A slim package fits in a card case to carry around.</p>																									
	<p>ACTIVE INGREDIENTS (in 6 troches) Cetylpyridinium Chloride Hydrate 8.28mg [Inactive ingredients] Hydroxypropylcellulose, Sucralose, Povidone, Pullulan, Macrogol, I-Menthol, Tannic Acid, D-Sorbitol, Saccharin Sodium, Sucrose Fatty Acid Ester, Flavor, New Coccine, Sunset Yellow FCF</p>	<p>For further information:</p> <table border="1"> <tr> <td>Manufactured and Distributed by:</td> <td colspan="3">TAIHO PHARMACEUTICAL CO., LTD.</td> </tr> <tr> <td>Dosage form</td> <td colspan="3">Troche</td> </tr> <tr> <td>Packaging unit</td> <td>12 troches</td> <td></td> <td></td> </tr> <tr> <td>Manufacturer's recommended price</td> <td>500 yen</td> <td></td> <td></td> </tr> <tr> <td>JAN Code</td> <td>45198618</td> <td></td> <td></td> </tr> <tr> <td>Expiration</td> <td colspan="3">3 years</td> </tr> </table> <p>DOSAGE AND ADMINISTRATION Put the following dose in the mouth, and slowly dissolve without chewing. Dose for Adults (15 years and over) 1 troche Frequency of daily dosing 4 to 6 times Do not use for children under 15 <Precautions> (1) Strictly follow the dosing instructions. (2) Do not chew or swallow whole.</p>	Manufactured and Distributed by:	TAIHO PHARMACEUTICAL CO., LTD.			Dosage form	Troche			Packaging unit	12 troches			Manufacturer's recommended price	500 yen			JAN Code	45198618			Expiration	3 years	
Manufactured and Distributed by:	TAIHO PHARMACEUTICAL CO., LTD.																								
Dosage form	Troche																								
Packaging unit	12 troches																								
Manufacturer's recommended price	500 yen																								
JAN Code	45198618																								
Expiration	3 years																								
<p>INDICATIONS Throat pain, throat swelling, sore throat, throat discomfort, and hoarse voice due to throat inflammation, sterilization of the mouth, and removal of bad breath</p>																									
<p>PRECAUTIONS FOR USE Consultation 1. The following persons should consult a physician, dentist or pharmacist before using the medicine. (1) Persons under treatment by a physician or dentist. (2) Pregnant women or possibly pregnant women. (3) The person who will take this drug or any on in the family has allergic constitution. (4) Persons with a history of allergic symptoms to any drug. 2. If any of these symptoms occur, discontinue the use of the medicine immediately, and consult a physician, dentist or pharmacist with this leaflet. (1) If any of the following symptoms occur. Skin: Rash, redness, itching (2) If symptoms do not improve after using the medicine for one week.</p>																									