Brand name: Solmack 5			
RISK CATEGORIES			
Designated quasi-drug			
Designated quasi-drug			
CHARACTERISTICS			
I Gastrointestinal medicine containing crude drugs			
The first product of Solmack series to be taken before			
meals			(707) EH-121
Easy-to-take refreshing plum flavor			
It contains a maximum of 2,000 mg of turmeric (maximum			
amount within the standard range for registered quasi-drug:	Man Carling And		
equivalent to the amount of crude drug per 50 mL bottle)	L'AST	JE COLLE	
The medicine contains perilla herb and red ginseng which	G	EJA	עיקעוע
help improve gastrointestinal function.	8	BRR. VA47-95	DELE
A designated quasi-drug "Solmack 5" contains five types of	本入り 50mL	定医薬部外品 一 大農業品工業株式生	ALL MARKAN
crude drugs.		http://www.taibs.co	JAN .
It exerts an excellent effect on stomach discomfort and			
feeling queasy due to overeating or overdrinking.			
ACTIVE INGREDIENTS	For further information:		
Per bottle (50 mL)		•	
Turmeric Extract 200 mg (equivalent to 2,000 mg of	Manufactured and		
turmeric)	Distributed by:		
Red Ginseng Extract 83.3 mg (equivalent to 333.2 mg of	Dosage form	e form Liquid	
red ginseng)			- -
Perilla Herb Extract 37 mg (equivalent to 333 mg of perilla	Packaging unit	50 mL x 1 bottle	50 mL x 8 bottles
herb) Clove Oil 5 mg	Manufacturer's	101/ 205	JPY 3,080
Dipotassium Glycyrrhizinate 7.2 mg	suggested retail price	JPY 385 (JPY 350 excluding tax)	(JPY 2,800 excluding tax)
Inactive ingredients: Glycerin, Fructose, Propylene Glycol,	JAN Code	45144646	4987117396010
Glucose, Povidone, Citric Acid Hydrate, Polyoxyethylene		0	
Hydrogenated Caster Oil, I-Menthol, Sodium Benzoate,	Expiration 3 years DOSAGE AND ADMINISTRATION		ears
pH Adjuster,Vanillin, Flavor			
• Shake well before use. As this medicine contains crud	Adults (15 years and over): Take 1 bottle (50 mL) before meals once daily.		
drugs, it may cause precipitation.			
		ren (under 15 years).	
	• Follow the dosing i	nstruction.	
	4		
INDICATIONS			
Stomach discomfort or feeling queasy (sickness, churning			
of stomach, hangover, drunken sickness, queasy, nausea)			
due to overeating or overdrinking			
PRECAUTIONS FOR USE	1		
1. The following persons should consult a physician or pharmacist before using the medicine.			
 Persons under treatment by a physician 			
2. In the following case, discontinue the use of the medicine, and consult a physician or pharmacist with this medicine.			
 If symptoms do not improve after using the medicine for 	a while		

CONTACT

For inquiries, please contact the shop where you purchased the product or the following: Manufacturer and distributor Contact:

Customer Relations Office, Taiho Pharmaceutical Co., Ltd. 1-27 Kandanishiki-cho, Chiyoda-ku, Tokyo, 101-8444, Japan

Phone: 0120-4527-66 Business hours: 9:00-17:00 (Excluding weekends and holidays) Website: <u>https://www.taiho.co.jp/</u>