Brand name: Solmack Digestive Drink Plus R

RISK CATEGORIES
Designated quasi-drug

CHARACTERISTICS
Stomach discomfort or feeling queasy (sickness, churning of stomach, hangover, drunken sickness, queasy, nausea)due to overeating or overdrinking



ACTIVE INGREDIENTS
Per bottle (50 mL)
Turmeric Extrac 30mg (equivalent to 300 mg of turmeric)
Isodonis herba Extract 15.46mg (equivalent to 201 mg ofisodonis herba)
Ginseng Extract 90.4 mg (equivalent to 606 mg of
Cinnamon Oil 1.5 mg
Clove oil 5 mg
Carnitine Hydrochloride 120 mg
Dipotassium Glycyrrhizinate7.2mg

Inactive ingredients: Fructose, Glucose, L-Menthol, Sodium Benzoate, Propylene Glycol, Polyoxyethylene Hydrogenated Castor Oil, pH Adjusting Agent, Benzyl Benzoate, Ethanol, Phenolic Compounds (Eugenol), Flavor(≥ 0.005 mL of alcohol)

- Shake well before use. As this medicine contains crud drugs, it may cause precipitation.

INDICATIONS
Stomach discomfort or feeling queasy (sickness, churning of stomach, hangover, drunken sickness, queasy, nausea) due to overeating or overdrinking

For further information:

Manufactured and Distributed by:	TAIHO PHARMACEUTICAL CO., LTD.	
Dosage form	Liquid	
Packaging unit	50 mL x 1 bottle	50 mL x 8 bottles
Manufacturer's suggested retail price	JPY 438 (JPY 398excluding)	JPY 3,502 (JPY 3,184 excluding)
JAN Code	45209680	4987117400427
Expiration	3 years	

DOSAGE AND ADMINISTRATION
Adults (15 years and over): Take 1 bottle (50 mL) before or between meals once daily.

- Do not use in children (under 15 years).
- Follow the dosing instruction.

PRECAUTIONS FOR USE
Consultation
1. In the following cases, discontinue the use of the medicine immediately, and consult a physician or pharmacist with this medicine.
(1) If rash occurs after using the medicine.
(2) If symptoms do not improve after using the medicine for a while.
2. Diarrhea may occur. If the symptom persists or gets worse, discontinue the use of the medicine, and consult a physician or pharmacist.

CONTACT

For inquiries, please contact the shop where you purchased the product or the following:
Manufacturer and distributor Contact:
Customer Relations Office, Taiho Pharmaceutical Co., Ltd. 1-27 Kandanishiki-cho, Chiyoda-ku, Tokyo, 101-8444, Japan
Phone: 0120-4527-66 Business hours: 9:00-17:00 (Excluding weekends and holidays)
Website: https://www.taiho.co.jp/