Brand name: Solmack Digestive Drink Plus

### RISK CATEGORIES

Designated quasi-drug

### CHARACTERISTICS

I Gastrointestinal medicine containing crude drugs
This medicine is effective for stomach discomfort and
felling queasy due to overdrinking and overeating.
It contains ginseng, turmeric, glycyrrhiza, isodonis herba,
mallotus bark, cinnamon bark, clove, and carnitine
hydrochloride.





#### ACTIVE INGREDIENTS

Per bottle (50 mL)

Turmeric Fluidextract 0.3 mL (equivalent to 300 mg of turmeric)

Glycyrrhiza Extract 10 mg (equivalent to 143 mg of glycyrrhiza)

Isodonis herba Extract 30 mg (equivalent to 201 mg of isodonis herba)

Ginseng Extract 200 mg (equivalent to 606 mg of ginseng) Mallotus Bark Extract 50 mg (equivalent to 500 mg of mallotus bark)

Cinnamon Oil 1.5 mg

Clove oil 5 mg

Carnitine Hydrochloride 120 mg

Inactive ingredients: Propylene Glycol, Fructose, Glucose, Alcohol, Polyoxyethylene Hardened Caster Oil, I-Menthol, Sodium Benzoate, pH Adjuster, Flavor (≥ 0.5 mL of alcohol)

• Shake well before use. As this medicine contains crud drugs, it may cause precipitation.

#### For further information:

Manufactured and Distributed by:	TAIHO PHARMACEUTICAL CO., LTD.	
Dosage form	Liquid	
Packaging unit	50 mL x 1 bottle	50 mL x 8 bottles
Manufacturer's suggested retail price	JPY 385 (JPY 350 excluding tax)	JPY 3,080 (JPY 2,800 excluding tax)
JAN Code	45144622	4987117398571
Expiration	3 years	

## DOSAGE AND ADMINISTRATION

Adults (15 years and over): Take 1 bottle (50 mL) before or between meals once daily.

- Do not use in children (under 15 years).
- Follow the dosing instruction.

# INDICATIONS

Stomach discomfort or feeling queasy (sickness, churning of stomach, hangover, drunken sickness, queasy, nausea) due to overeating or overdrinking

### PRECAUTIONS FOR USE

# Consultation

- 1. In the following cases, discontinue the use of the medicine immediately, and consult a physician or pharmacist with this medicine.
- (1) If rash occurs after using the medicine.
- (2) If symptoms do not improve after using the medicine for a while.
- 2. Diarrhea may occur. If the symptom persists or gets worse, discontinue the use of the medicine, and consult a physician or pharmacist.