


Brand name: Solmack EX 2																				
<p>RISK CATEGORIES Second-class OTC drug</p> <p>CHARACTERISTICS</p> <ul style="list-style-type: none"> Gastrointestinal medicine containing crude drugs <p>Nine types of crude drugs are effective. (Isodonis Herba, Glycyrrhiza, Gentian, Coptis Rhizome, Cinnamon Bark, Turmeric, Clove, Ginseng, Atractylodes Lancea Rhizome) + Carnitine Hydrochloride This medicine contains turmeric and ginseng, and effective for overeating (hyperphagia), overdrinking (binge drinking), heavy stomach, loss of appetite (decreased appetite), and dysplasia.</p>																				
	<p>ACTIVE INGREDIENTS</p> <p>- Per bottle (50 mL)- Quantities described in parenthesis are those equivalent to relevant crude drugs.</p> <p>Isodonis Herba Extract 30 mg (201 mg) Turmeric Fluidextract 0.3 mL (300 mg) Glycyrrhiza Extract 115 mg (1,644.5 mg) Clove Tincture 0.126 mL (37.8 mg) Gentian Tincture 0.376 mL (112.8 mg) Ginseng Fluidextract 0.5 mL (500 mg) Coptis Rhizome Tincture 0.792 mL (237.6 mg) Atractylodes Lancea Rhizome Fluidextract 1.2 mL (1,200 mg) Cinnamon Tincture 0.25 mL (75 mg) Carnitine Hydrochloride 120 mg</p> <p>Inactive ingredients: White Soft Sugar, Honey, Alcohol, Propylene Glycol, Acacia, Edetate calcium disodium, l-Menthol, Sodium Benzoate, pH Adjuster, Flavor (≤1.5 mL of alcohol)</p> <ul style="list-style-type: none"> Shake well before use. As this medicine contains crud drugs, it may cause precipitation. 																			
<p>For further information:</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 33%;">Manufactured and Distributed by:</td> <td colspan="2">TAIHO PHARMACEUTICAL CO., LTD.</td> </tr> <tr> <td>Dosage form</td> <td colspan="2" style="text-align: center;">Liquid</td> </tr> <tr> <td>Packaging unit</td> <td style="text-align: center;">50 mL x 1 bottle</td> <td style="text-align: center;">50 mL x 8 bottles</td> </tr> <tr> <td>Manufacturer's suggested retail price</td> <td style="text-align: center;">JPY 638 (JPY 580 excluding tax)</td> <td style="text-align: center;">JPY 5,104 (JPY 4,640 excluding tax)</td> </tr> <tr> <td>JAN Code</td> <td style="text-align: center;">45144639</td> <td style="text-align: center;">4987117397895</td> </tr> <tr> <td>Expiration</td> <td colspan="2" style="text-align: center;">3 years</td> </tr> </table>			Manufactured and Distributed by:	TAIHO PHARMACEUTICAL CO., LTD.		Dosage form	Liquid		Packaging unit	50 mL x 1 bottle	50 mL x 8 bottles	Manufacturer's suggested retail price	JPY 638 (JPY 580 excluding tax)	JPY 5,104 (JPY 4,640 excluding tax)	JAN Code	45144639	4987117397895	Expiration	3 years	
Manufactured and Distributed by:	TAIHO PHARMACEUTICAL CO., LTD.																			
Dosage form	Liquid																			
Packaging unit	50 mL x 1 bottle	50 mL x 8 bottles																		
Manufacturer's suggested retail price	JPY 638 (JPY 580 excluding tax)	JPY 5,104 (JPY 4,640 excluding tax)																		
JAN Code	45144639	4987117397895																		
Expiration	3 years																			
<p>INDICATIONS</p> <p>Overeating (hyperphagia), overdrinking (binge drinking), heavy stomach, felling queasy (hangover, drunken sickness, sickness, churning of stomach, queasy, nausea), vomiting, weak stomach, loss of appetite (decreased appetite), stomach/abdominal bloating, dyspepsia, heartburn, heavy feeling in the chest</p>																				
<p>PRECAUTIONS FOR USE</p> <p><input checked="" type="checkbox"/> The following precaution should be observed. (If not observed, it may aggravate the current symptoms or increase the risk of adverse reactions.)</p> <ul style="list-style-type: none"> Do not use for a long period of time. <p>Consultation</p> <p>1. The following persons should consult a physician, pharmacist or registered salesperson before using the medicine.</p> <ol style="list-style-type: none"> (1) Persons under treatment by a physician (2) Elderly (3) Persons with oedema 																				
<p>DOSAGE AND ADMINISTRATION</p> <p>Adults (15 years and over): Take 1 bottle (50 mL) once daily.</p> <ul style="list-style-type: none"> Do not use in children (under 15 years). Strictly follow the dosing instruction. 																				

(4) Persons diagnosed with hypertension, kidney disease or cardiac disease

2. The following serious symptoms may rarely occur after using the medicine. The following symptoms may be adverse reactions to the medicine. If any of these symptoms occur, discontinue the use of the medicine immediately, and consult a physician with this medicine.
Pseudoaldosteronism, myopathy (Weakness, numbness, tightness, and stiffness of limbs occur along with feelings of weakness and muscle pain, and they gradually get worse.)
3. If symptoms do not improve after using the medicine for five to six days, discontinue the use of the medicine, and consult a physician, pharmacist or registered salesperson with this medicine.

CONTACT

For inquiries, please contact the shop where you purchased the product or the following:

Manufacturer and distributor Contact:

Customer Relations Office, Taiho Pharmaceutical Co., Ltd. 1-27 Kandanshiki-cho, Chiyoda-ku, Tokyo, 101-8444, Japan

Phone: 0120-4527-66 Business hours: 9:00-17:00 (Excluding weekends and holidays)

Website: <https://www.taiho.co.jp/>