Brand name: Solmack Gold

RISK CATEGORIES

Designated quasi-drug

CHARACTERISTICS

I Gastrointestinal medicine containing crude drugs Ten types of crude drugs such as turmeric and glycyrrhiza are effective.

(Turmeric, Glycyrrhiza, Isodonis Herba, Cardamon, Citri Unshiu Pericarpium Immaturus, Acori Graminei Rhizoma, Saussurea Root, Mallotus Bark, Cinnamon Bark, Clove) + Carnitine Hydrochloride

This medicine is effective for stomach discomfort or feeling queasy (sickness, churning of stomach, hangover, drunken sickness, queasy, nausea) due to overeating or overdrinking.





ACTIVE INGREDIENTS

Per bottle (50 mL)

Turmeric Fluidextract 0.6 mL (equivalent to 600 mg of turmeric)

Glycyrrhiza Extract 11.6 mg (equivalent to 166 mg of glycyrrhiza)

Isodonis Herba Extract 90 mg (equivalent to 603 mg of isodonis herba)

Cardamon Fluidextract 0.6 mL (equivalent to 600 mg of cardamom)

Citri Unshiu Pericarpium Immaturus 150 mg (equivalent to 600 mg of citri unshiu pericarpium immaturus)

Mallotus Bark Extract 90 mg (equivalent to 603 mg of mallotus bark)

Saussurea Root Extract 129 mg (equivalent to 993 mg of saussurea root)

Mallotus Bark Extract 50 mg (equivalent to 500 mg of mallotus bark)

Cinnamon Oil 3.0 mg

Clove Oil 6.6 mg

Carnitine Hydrochloride 120 mg

Inactive ingredients: Propylene Glycol, Fructose, Glucose, Alcohol, Polyoxyethylene Hardened Caster Oil, I-Menthol, Sodium Benzoate, pH Adjuster, Flavor (≥ 0.5 mL of alcohol)

• Shake well before use. As this medicine contains crud drugs, it may cause precipitation.

For further information:

Manufactured and Distributed by:	TAIHO PHARMACEUTICAL CO., LTD.	
Dosage form	Liquid	
Packaging unit	50 mL x 1 bottle	50 mL x 8 bottles
Manufacturer's suggested retail price	JPY 523 (JPY 475 excluding tax)	JPY 4,180 (JPY 3,800 excluding tax)
JAN Code	4987117398717	4987117398724
Expiration	3 years	

DOSAGE AND ADMINISTRATION

Adults (15 years and over): Take 1 bottle (50 mL) before or between meals once daily.

- Do not use in children (under 15 years).
- Follow the dosing instruction.

INDICATIONS

Stomach discomfort or feeling queasy (sickness, churning of stomach, hangover, drunken sickness, queasy, nausea) due to overeating or overdrinking

PRECAUTIONS FOR USE

- 1. The following persons should consult a physician or pharmacist before using the medicine.
 - Persons under treatment by a physician
- 2. In the following case, discontinue the use of the medicine, and consult a physician or pharmacist with this medicine.
 - If symptoms do not improve after using the medicine for a while