Brand name: Solmack Plus

RISK CATEGORIES

Second-class OTC drug

CHARACTERISTICS

Gastrointestinal medicine containing crude drugs

Eight types of crude drugs are effective.
(Turmeric, Glycyrrhiza, Ginseng, Coptis
Rhizome, Clove, Gentian, Atractylodes Lancea
Rhizome, Cinnamon Bark) + Carnitine
Hydrochloride

The medicine is effective for felling queasy (hangover, drunken sickness, sickness, churning of stomach, queasy, nausea), vomiting, dyspepsia, overeating (hyperphagia), overdrinking (binge drinking), and loss of appetite.

It contains turmeric and glycyrrhiza, and has a cool menthol taste.







ACTIVE INGREDIENTS

Per 2 bottles (50 mL) daily

Turmeric Fluidextract 0.3 mL (equivalent to 300 mg of the crude drug)

Glycyrrhiza Extract 135 mg (equivalent to 1,930.5 mg of the crude drug)

Ginseng Fluidextract 0.5 mL (equivalent to 500 mg of the crude drug)

Coptis Rhizome Tincture 0.826 mL (equivalent to 247.8 mg of the crude drug)

Clove Tincture 0.126 mL (equivalent to 37.8 mg of clove)

Gentian Tincture 0.376 mL (equivalent to 112.8 mg of the crude drug)

Atractylodes Lancea Rhizome Fluidextract 1.2 mL (equivalent to 1,200 mg of the crude drug) Cinnamon Bark Tincture 0.25 mL (equivalent to 75 mg of the crude drug) Carnitine Hydrochloride 120 mg

Inactive ingredients: Honey, White Soft Sugar, Acacia, Edetate calcium disodium, *I*-Menthol, Alcohol, Sodium Benzoate, pH Adjuster, Benzyl Benzoate, Flavor (≤0.8mL of alcohol/bottle)

 Shake well before use. As this medicine contains crud drugs, it may cause precipitation.

INDICATIONS

Felling queasy (hangover, drunken sickness, sickness, churning of stomach, queasy, nausea), vomiting, dyspepsia, overeating (hyperphagia), overdrinking (binge drinking), heavy stomach, weak stomach, loss of appetite (decreased appetite), stomach/abdominal bloating, heartburn, heavy feeling in the chest

For further information:

Manufactured and Distributed by:	TAIHO PHARMACEUTICAL CO., LTD.		
Dosage form	Liquid		
Packaging unit	25 mL x 1 bottle	25 mL x 2 bottles	25 mL x 5 bottles
Manufacturer's suggested retail price	JPY 440 (JPY 400 excluding tax)	JPY 880 (JPY 800 excluding tax)	JPY 2,200 (JPY 2,000 excluding tax)
JAN Code	45144608	4987117398816	4987117398847
Expiration	3 years		

DOSAGE AND ADMINISTRATION

Adults (15 years and over): Take 1 bottle (25 mL) twice daily. Each dose should be taken at an interval of more than 4 hours.

- Do not use in children (under 15 years).
- Strictly follow the dosing instruction.

PRECAUTIONS FOR USE

Make The following precaution should be observed.

(If not observed, it may aggravate the current symptoms or increase the risk of adverse reactions.)

Do not use for a long period of time.

Consultation

- 1. The following persons should consult a physician, pharmacist or registered salesperson before using the medicine.
 - (1) Persons under treatment by a physician
 - (2) Elderly
 - (3) Persons with oedema
 - (4) Persons diagnosed with hypertension, kidney disease or cardiac disease
- 2. The following serious symptoms may rarely occur after using the medicine. The following symptoms may be adversereactions to the medicine. If any of these symptoms occur, discontinue the use of the medicine immediately, and consult a physician with this medicine.
 - Pseudoaldosteronism, myopathy (Weakness, numbness, tightness, and stiffness of limbs occur along with feelings of weakness and muscle pain, and they gradually get worse.)
- 3. If symptoms do not improve after using the medicine for five to six days, discontinue the use of the medicine, and consulta physician, pharmacist or registered salesperson with this medicine.

CONTACT

For inquiries, please contact the shop where you purchased the product or the following: Manufacturer and distributor Contact:

Customer Relations Office, Taiho Pharmaceutical Co., Ltd. 1-27 Kandanishiki-cho, Chiyoda-ku, Tokyo, 101-8444, Japan Phone: 0120-4527-66 Business hours: 9:00-17:00 (Excluding weekends and holidays)

Website: https://www.taiho.co.jp/