

Brand name: Solmack Q-Arl S

RISK CATEGORIES
Designated quasi-drug
Vitamin-containing health drug

CHARACTERISTICS

- L-cysteine supports alcohol metabolism, which may help prevent tiredness associated with hangovers.
- Taking vitamins may help skin problems such as rough skin and dry skin.



ACTIVE INGREDIENTS: In daily dose (2 tablets)
Ascorbic Acid (Vitamin C) 300 mg
L-Cysteine 80 mg
D-Alpha Tocopheryl Succinate (Vitamin E Natural) 20 mg
Riboflavin (Vitamin B₂) 2 mg
Pyridoxine Hydrochloride (Vitamin B₆) 4 mg

Inactive ingredients: Corn Starch, Lactose, Cellulose, Carmellose, Hydroxypropylcellulose, Magnesium Stearate, Hypromellose, Titanium Oxide, Talc, Macrogol, Carnauba Wax

<Precautions of ingredients and amount>

- 1) This drug may interfere with urine and stool tests. If you are given such a test, inform your doctor that you are taking this drug.
- 2) You may experience yellow urine after taking this drug, which is due to riboflavin (vitamin B₂) and is harmless.

INDICATIONS

- Improvement and prevention of poor health associated with malnutrition in daily life
 - ◆ Loss of appetite and tiredness associated with hangovers
 - ◆ Skin problems such as rough skin and dry skin

For further information:

Distributor	TAIHO PHARMACEUTICAL CO., LTD.		
Dosage form	Tablets		
Packaging unit	2 doses (4 tablets)	10 doses (20 tablets)	
Manufacturer's suggested retail price	JPY 330 (JPY 300 excluding tax)	JPY 1,628 (JPY 1,480 excluding tax)	
JAN code	4987117400014	4987117400212	
Expiration	3 years		

DOSAGE AND ADMINISTRATION

Take the following doses with cold or warm water.

Adult (15 years old and over)	2 tablets per dose	Once daily
Children under 15 years old	Do not take this product.	

<Precautions for dosage and administration>

Follow the dosage and administration instructions. (If other products including vitamins are taken concomitantly, be careful of overdose.)

PRECAUTIONS FOR USE AND CONSULTATIONS

1. If the following symptoms are observed after taking this medicine, they may be adverse reactions, so immediately discontinue use of this medicine and consult a physician, pharmacist, or registered salesperson, while showing them the product.

Affected part	Symptoms
Skin	Rash/redness, itching
Gastrointestinal system	Nausea/vomiting, feeling of discomfort in the stomach

2. The following symptoms may be observed after taking this medicine. If these symptoms persist or worsen, discontinue use of this medicine and consult a physician, pharmacist, or registered salesperson, while showing them the product. Diarrhea, constipation
3. When symptoms do not improve even after taking this medicine for a while, stop taking this medicine and consult a physician, pharmacist, or registered salesperson, while showing them the product.

PRECAUTIONS FOR STORAGE AND HANDLING

[4 tablets]

- (1) Store in a dry and cool place, away from direct sunlight.
- (2) Keep out of the reach of children.
- (3) Do not transfer to other containers. (It may cause misuse or change the quality)
- (4) Do not use the product past the expiration date.

[20 tablets]

- (1) Tightly seal the zipper bag and store in a dry and cool place, away from direct sunlight.
- (2) Keep out of the reach of children.
- (3) Do not transfer to other containers. (It may cause misuse or change the quality)
- (4) Do not use the product past the expiration date.
- (5) Do not touch the product with wet hands. If tablets are returned to the bag after being touched by wet hands, they may affect the other tablets.
- (6) Keep the desiccant inside the bag until all tablets are taken. Also, be careful not to ingest the desiccant accidentally.

CONTACT

For inquiries, please contact the shop where you purchased the product or the following:

Distributor

Contact: Customer Relations Office, Taiho Pharmaceutical Co., Ltd.

1-27 Kandnishiki-cho, Chiyoda-ku, Tokyo, 101-8444, Japan

Phone: 0120-4527-66

Business hours: 9:00-17:00 (Excluding weekends and holidays)

Website: <https://www.taiho.co.jp/>

Manufacturer and distributor

Kyoto Pharmaceutical Industries, Ltd.

38 Nishinokyo Tsukinowa-cho, Nakagyo-ku, Kyoto 604-8444, Japan