Brand name: Tiovita Drink 1000

# RISK CATEGORIES

Third-class OTC drug

#### **CHARACTERISTICS**

I Nutrient tonic drink

- Third-class OTC drug Red Tiovita!
- It contains 1,000 mg of taurine as an active ingredient.
- It contains vitamin B1 and vitamin B6 twice and 10 times more than the conventional product (Tiovita Drink), respectively.
- Easy-to-drink "mixed fruit taste"
- It only has "68 kcal".



### **ACTIVE INGREDIENTS**

Per 100 mL bottle

Taurine 1,000 mg, Inositol 50 mL, Nicotinamide 20 mg, Thiamine Nitrate (Vitamin B1) 10 mg, Riboflavin Sodium Phosphate (Vitamin B2) 5 mg, Pyridoxine Hydrochloride (Vitamin B6) 50 mg, Carnitine Hydrochloride 100 mg, Anhydrous Caffeine 50 mg

Inactive ingredients: White Soft Sugar, Glucose, DL-Malic Acid, Propyl Gallate, Sodium Benzoate, pH Adjuster, Vanillin, Ethyl Vanillin, Flavor

 Urine may turn yellow because of vitamin B2 contained in the medicine.
It is nothing to worry about.

## **INDICATIONS**

Nutritional fortification, weak constitution, and supply of nutrients in the following cases: physical fatigue, during and after illness, loss of appetite, nutritional disorders, febrile debilitating illness, during pregnancy or lactation

### For further information:

Manufactured and Distributed by:	TAIHO PHARMACEUTICAL CO., LTD.		
Dosage form	Liquid		
Packaging unit	100 mL x 1 bottle	100 mL x 10 bottles	100 mL x 50 bottles
Manufacturer's suggested retail price	JPY 176 (JPY 160 excluding tax)	JPY 1,760 (JPY 1,600 excluding tax)	JPY 8,800 (JPY 8,000 excluding tax)
JAN Code	4987117437805	4987117437812	4987117437829
Expiration	3 years		

### DOSAGE AND ADMINISTRATION

Adults (15 years and over): Take 1 bottle (100 mL) once daily.

- Do not use in children (under 15 years).
- Follow the dosing instruction.

### PRECAUTIONS FOR USE

- Consultation
- 1. Skin rash may be adverse reactions to the medicine. If the symptom occurs, discontinue the use of the medicineimmediately, and consult a physician, pharmacist or registered salesperson with this medicine.
- 2. Diarrhea may occur after using the medicine. If the symptom persists or gets worse, discontinue the use of the medicine, and consult a physician, pharmacist or registered salesperson with this medicine.
- 3. If symptoms do not improve after using the medicine for a while, discontinue the use of the medicine, and consult aphysician, pharmacist or registered salesperson with this medicine.