Brand name: Tiovita Drink 2000

# RISK CATEGORIES

Designated quasi-drug

# CHARACTERISTICS

I Vitamin health drink

- "Effective" and "Good Taste" Green Tiovita! with a catchphrase "One Love to Share"!
- It contains 2,000 mg of taurine as an active ingredient (twice more than Tiovita Drink).
- Easy-to-take "cassis flavor"
- It only has "72 kcal".







# **ACTIVE INGREDIENTS**

Per 100 mL bottle

Taurine 2,000 mg, Inositol 50 mg, Nicotinamide 20 mg, Thiamine Nitrate (Vitamin B1) 5 mg, Riboflavin Sodium Phosphate (Vitamin B2) 5 mg, Pyridoxine Hydrochloride (Vitamin B6) 5 mg, Carnitine Hydrochloride 100 mg, Anhydrous Caffeine 50 mg

Inactive ingredients: White Soft Sugar, Fructose, Glucose, DL-Malic Acid, Propyl Gallate, Sodium Benzoate, pH Adjuster, Vanillin, Flavor

 Urine may turn yellow because of vitamin B2 contained in the medicine. It is nothing to worry about.

#### For further information:

Manufactured and Distributed by:	TAIHO PHARMACEUTICAL CO., LTD.			
Dosage form	Liquid			
Packaging unit	100 mL x 1 bottle	100 mL x 3 bottles	100 mL x 10 bottles	100 mL x 50 bottles
Manufacturer's suggested retail price	JPY 198 (JPY 180 excluding tax)	JPY 594 (JPY 540 excluding tax)	JPY 1,980 (JPY 1,800 excluding tax)	JPY 9,900 (JPY 9,000 excluding tax)
JAN Code	4987117439106	4987117439205	4987117439113	4987117439120
Expiration	3 years			

# DOSAGE AND ADMINISTRATION

Adults (15 years and over): Take 1 bottle (100 mL) once daily.

- Do not use in children (under 15 years).
- Follow the dosing instruction. (Please pay attention to overdose if other products containing vitamins are concomitantly used.)

# INDICATIONS

ONutritional fortification
OWeak constitution
OSupply of nutrients in the following
cases:, physical fatigue, during and
after illness, loss of appetite, nutritional
disorders, febrile debilitating illness,
during pregnancy or lactation

# PRECAUTIONS FOR USE

- Consultation
- The following cases, discontinue the use of the medicine immediately, and consult a physician or pharmacist with this medicine.
- (1) If rash occurs after using the medicine.
- (2) If symptoms do not improve after using the medicine for a while.
- 2. Diarrhea may occur. If the symptom persists or gets worse, discontinue the use of the medicine, and consult a physician or pharmacist.