


Brand name: Tiovita Drink Aibitasu				
RISK CATEGORIES Third-class OTC drug CHARACTERISTICS I Nutrient tonic drink - "Blueberry flavor" which is easy to take for women - The medicine contains vitamin B1, vitamin B2, and vitamin B6 twice, three times, and 10 times more than the conventional product (Tiovita Drink), respectively. - It contains 1,000 mg of taurine as an active ingredient. - It only has "68 kcal".				
				
ACTIVE INGREDIENTS Per 100 mL bottle Taurine 1,000 mg, Inositol 50 mg, Nicotinamide 20 mg, Thiamine Nitrate (Vitamin B1) 10 mg, Riboflavin Sodium Phosphate (Vitamin B2) 15 mg, Pyridoxine Hydrochloride (Vitamin B6) 50 mg, Carnitine Hydrochloride 120 mg, Anhydrous Caffeine 50 mg Inactive ingredients: White Soft Sugar, Glucose, DL-Malic Acid, Propyl Gallate, Sodium Benzoate, pH Adjuster, Vanillin, Ethyl Vanillin, Flavor <ul style="list-style-type: none"> Urine may turn yellow because of vitamin B2 contained in the medicine. It is nothing to worry about. 				
INDICATIONS Nutritional fortification, weak constitution, and supply of nutrients in the following cases: physical fatigue, during and after illness, loss of appetite, nutritional disorders, febrile debilitating illness, during pregnancy or lactation				
For further information:				
Manufactured and Distributed by:		TAIHO PHARMACEUTICAL CO., LTD.		
Dosage form		Liquid		
Packaging unit	100 mL x 1 bottle	100 mL x 3 bottles	100 mL x 10 bottles	100 mL x 50 bottles
Manufacturer's suggested retail price	JPY 198 (JPY 180 excluding tax)	JPY 594 (JPY 540 excluding tax)	JPY 1,980 (JPY 1,800 excluding tax)	JPY 9,900 (JPY 9,000 excluding tax)
JAN Code	4987117439304	4987117439335	4987117439311	4987117439328
Expiration	3 years			
DOSAGE AND ADMINISTRATION Adults (15 years and over): Take 1 bottle (100 mL) once daily. <ul style="list-style-type: none"> Do not use in children (under 15 years). Follow the dosing instruction. 				

PRECAUTIONS FOR USE

Consultation

1. Skin rash may be adverse reactions to the medicine. If the symptom occurs, discontinue the use of the medicine immediately, and consult a physician, pharmacist or registered salesperson with this medicine.
2. Diarrhea may occur after using the medicine. If the symptom persists or gets worse, discontinue the use of the medicine, and consult a physician, pharmacist or registered salesperson with this medicine.
3. If symptoms do not improve after using the medicine for a while, discontinue the use of the medicine, and consult a physician, pharmacist or registered salesperson with this medicine.