

Brand name: Tiovita Drink

RISK CATEGORIES
Designated quasi-drug

CHARACTERISTICS
I Vitamin health drink

- Long-time beloved vitamin healthdrink with a catchphrase "One Love to Share to help relieve physical exhaustion"
- It contains 1,000 mg of taurine, vitamin B groups, and carnitine hydrochloride as active ingredients.
- Easy-to-take "mixed fruit flavor"
- It only has "70 kcal".



ACTIVE INGREDIENTS
Per 100 mL bottle
Taurine 1,000 mg, Inositol 50 mg, Nicotinamide 20 mg, Thiamine Nitrate (Vitamin B1) 5 mg, Riboflavin Sodium Phosphate (Vitamin B2) 5 mg, Pyridoxine Hydrochloride (Vitamin B6) 5 mg, Carnitine Hydrochloride 100 mg, Anhydrous Caffeine 50 mg

Inactive ingredients: High-fructose Corn Syrup, DL-Malic Acid, Propyl Gallate, Sodium Benzoate, pH Adjuster, Citric Acid, Sodium Citrate, D-Sorbitol, Vanillin, Ethyl Vanillin, Flavor

- Urine may turn yellow because of vitamin B2 contained in the medicine. It is nothing to worry about.

For further information:

Manufactured and Distributed by:	TAIHO PHARMACEUTICAL CO., LTD.				
Dosage form	Liquid				
Packaging unit	100 mL x 1 bottle	100 mL x 2 bottles	100 mL x 3 bottles	100 mL x 10 bottles	100 mL x 50 bottles
Manufacturer's suggested retail price	JPY 161 (JPY 146 excluding tax)	JPY 321 (JPY 292 excluding tax)	JPY 482 (JPY 438 excluding tax)	JPY 1,606 (JPY 1,460 excluding tax)	JPY 8,030 (JPY 7,300 excluding tax)
JAN Code	4987117438710	4987117438956	4987117438918	4987117438741	4987117438802
Expiration	3 years				

INDICATIONS

- Recovery from and prophylaxis of fatigue
- Improvement and prophylaxis of physical malfunction associated with malnutrition in daily living
◇Fatiguability, persistent fatigue, lack of physical strength, heaviness in the body, feeling tired
- Maintenance and improvement of physical strength, resistance, or concentration
- Supply of nutrients in the following cases: decreased physical strength during/after illness, febrile debilitating illness, loss of appetite, during pregnancy/lactation or before/after delivery

DOSAGE AND ADMINISTRATION
Adults (15 years and over): Take 1 bottle (100 mL) once daily.

- Do not use in children (under 15 years).
- Follow the dosing instruction. (Please pay attention to overdose if other products containing vitamins are concomitantly used.)

PRECAUTIONS FOR USE

Consultation

1. If skin rash occurs after using the medicine, this may be an adverse reaction of the medicine. Discontinue the use of the medicine immediately, and consult a physician, pharmacist, or registered salesperson with this medicine.
2. Diarrhea may occur after using the medicine. If the symptom persists or gets worse, discontinue the use of the medicine, and consult a physician, pharmacist, or registered salesperson with this medicine.
3. If symptoms do not improve after using the medicine for a while, discontinue the use of the medicine, and consult a physician, pharmacist, or registered salesperson with this medicine.