

Brand name: Tiovita Gold			
RISK CATEGORIES Second-class OTC drug CHARACTERISTICS I Nutrient tonic drink - The medicine contains royal jelly, ginseng, and epimedium herb which are effective for severe fatigue! (For further information, please see ACTIVE INGREDIENTS.) - It is also good for supply of nutrients at the time of common cold! - Easy-to-take 30 mL drink - Sweat and mild "carrot flavor" - It only has "34 kcal".			
	ACTIVE INGREDIENTS Per 30 mL bottle Royal Jelly 100 mg, Ginseng Fluidextract 0.6 mL (equivalent to 600 mg of ginseng), Epimedium Herb Fluidextract 0.1 mL (equivalent to 100 mg of epimedium herb), Thiamine Nitrate (Vitamin B1) 5 mg, Riboflavin Sodium Phosphate (Vitamin B2) 5 mg, Pyridoxine Hydrochloride (Vitamin B6) 5 mg, Nicotinamide 30 mg, Anhydrous Caffeine 50 mg (≤ 0.2 mL of alcohol) Inactive ingredients: Citric Acid, , Propyl Gallate, Alcohol, Sodium Benzoate, pH Adjuster, Vanillin, Corn Starch, Flavor (1) Shake well before use. As this medicine contains crude drugs, it may cause precipitation. (2) Urine may turn yellow because of vitamin B2 contained in the medicine. It is nothing to worry about.		
For further information:			
Manufactured and Distributed by:		TAIHO PHARMACEUTICAL CO., LTD.	
Dosage form		Liquid	
Packaging unit	30 mL x 1 bottle	30 mL x 3 bottles	30 mL x 10 bottles
Manufacturer's suggested retail price	JPY 385 (JPY 350 excluding tax)	JPY 1,155 (JPY 1,050 excluding tax)	JPY 3,850 (JPY 3,500 excluding tax)
JAN Code	49684315	4987117437218	4987117437355
Expiration		3 years	
DOSAGE AND ADMINISTRATION Adults (15 years and over): Take 1 bottle (30 mL) once daily. ● Do not use in children (under 15 years). ● Follow the dosing instruction.			
INDICATIONS Supply of nutrients in the following cases: febrile debilitating illness, physical fatigue, during and after illness, loss of appetite, nutritional disorders, during pregnancy or lactation, and nutritional fortification, weak constitution			
PRECAUTIONS FOR USE ■ Consultation 1. Diarrhea may occur after using the medicine. If the symptom persists or gets worse, discontinue the use of the medicine, and consult a physician, pharmacist or registered salesperson with this medicine. 2. If symptoms do not improve after using the medicine for a while, discontinue the use of the medicine, and consult a physician, pharmacist or registered salesperson with this medicine.			