Brand name: Urepearl Plus Cream

# RISK CATEGORIES Second-class OTC drug

### **CHARACTERISTICS**

I Medicine for the treatment of dry skin

- This medicine contains a natural moisturizing factor, urea and vitamin E (tocopheryl acetate) which help to maintain water, and smoothen and moisturize dry skin.
- It contains three types of components as anti-itching active ingredients.
- It is a cream which fits to the skin and spreads smoothly.





### **ACTIVE INGREDIENTS**

**INDICATIONS** 

This is a cream containing the following active ingredients per 100 g.

Urea 10.0 g, Diphenhydramine Hydrochloride 1.0 g, d-Camphor 1.0 g, Lidocaine 2.0 g, Tocopherol Acetate 0.3 g Inactive Ingredients: Methyl Paraben, Butylated Hydroxytoluene (BHT), Glyceryl Triisooctanoate, Octyldodecyl Myristate, Cholesterol, Methyl Polysiloxane, Stearyl Alcohol, Cetanol, Behenyl Alcohol, Glyceryl Stearate, Sorbitan Stearate, Polyoxyethylene Behenyl Ether, Macrogol Stearate, Polyoxyethylene Cetylether Sodium Phosphate, Glycine, DL-Alanine, Sodium Chloride, Glycerin, Sodium Lactate, Lactic Acid, Purified Water

### For further information:

Distributed by:	TAIHO PHARMACEUTICAL CO., LTD.
Manufactured and Distributed by:	Otsuka Pharmaceutical Factory Inc.
Dosage form	Cream
Packaging unit	80 g
Manufacturer's suggested retail price	JPY 1,760 (JPY 1,600 excluding tax)
JAN Code	4987117140125
Expiration	3 years

### Apply an a

Dry skin accompanied by itching (xeroderma in the elderly/adults, dry skin in children)

# DOSAGE AND ADMINISTRATION

Apply an appropriate amount to the affected area once to several times daily.

### <Pre><Pre>cautions>

- (1) Be careful not to get the medicine into the eyes. If the medicine gets into the eyes, immediately wash with water or lukewarm water. If symptoms are severe, consult an ophthalmologist.
- (2) For children, use under parental guidance.
- (3) For external use only.
- (4) Clean the affected area and surrounding skin before use.
- (5) Follow the prescribed dosing instructions.
- (6) As this is not a cosmetics, use only for the affected areas listed in the indications, and do not use on the face for the purpose of basic skin care, etc.

### PRECAUTIONS FOR USE

∑ The following precaution should be observed.

(If not observed, it may aggravate the current symptoms or increase the risk of adverse reactions.) Do not apply to the following areas.

- (1) Around the eyes, mucous membranes (e.g. mouth, nasal cavity, vagina, etc.)
- (2) Inflammation or wounds
- (3) Severe skin sore or cracking of the skin



# Consultation

- 1. The following persons should consult a physician, pharmacist or registered distributor before using the medicine.
- (1) Persons under treatment by a physician
- (2) Persons with a history of allergic symptoms (e.g. rash, redness, itching, contact dermatitis, etc.) or irritation symptoms

# due to any drug

2. The following symptoms may be adverse reactions to the medicine. If any of these symptoms occur, discontinue the use of the medicine immediately, and consult a physician, pharmacist or registered distributor with this leaflet.

Affected body system	Symptoms
	Contact dermatitis, rash, redness, itching, irritation (pain, feeling hot, tingling sensation), swelling, scab-like skin exfoliation

3. If symptoms do not improve after using the medicine for about two weeks, discontinue the use of the medicine, and consult a physician, pharmacist or registered distributor with this leaflet.

# CONTACT

For inquiries, please contact the shop where you purchased the product or the following:

Distributor Contact: Customer Relations Office, Taiho Pharmaceutical Co., Ltd.

1-27 Kandanishiki-cho, Chiyoda-ku, Tokyo, 101-8444, Japan

Phone: 0120-4527-66 Business hours: 9:00-17:00 (Excluding weekends and holidays)

Website: https://www.taiho.co.jp/

Manufactured and Distributed : Otsuka Pharmaceutical Factory Inc.