

Brand name: Urepearl Plus Lotion 10

**RISK CATEGORIES**  
Second-class OTC drug

**CHARACTERISTICS**

I Medicine for the treatment of dry skin

- This medicine contains a natural moisturizing factor, urea and vitamin E (tocopheryl acetate) which help to maintain water, and smoothen and moisturize dry skin.
- It contains two types of components as anti-itching active ingredients.
- It is a clear and colorless lotion which spreads smoothly.



**ACTIVE INGREDIENTS**

Each 100 mL contains the following active ingredients.

Urea 10.0 g, Lidocaine 2.0 g, Diphenhydramine Hydrochloride 1.0 g

Inactive ingredients: Glycerin, Diisopropyl Sebacate, Lactic Acid, Sodium Lactate, Hydroxypropylcellulose, Purified Water, Flavor, Ethanol

For further information:

Distributed by:	TAIHO PHARMACEUTICAL CO., LTD.
Manufactured and Distributed by:	Otsuka Pharmaceutical Factory Inc.
Dosage form	Lotion
Packaging unit	100 mL
Manufacturer's suggested retail price	JPY 1,760 (JPY 1,600 excluding tax)
JAN Code	4987117140323
Expiration	3 years

**DOSAGE AND ADMINISTRATION**

Apply an appropriate amount to the affected area once to several times daily.

<Precautions>

- (1) Be careful not to get the medicine into the eyes. If the medicine gets into the eyes, immediately wash with water or lukewarm water. If symptoms are severe, consult an ophthalmologist.
- (2) For children, use under parental guidance.
- (3) For external use only.
- (4) Clean the affected area and surrounding skin before use.
- (5) Follow the prescribed dosing instructions.
- (6) As this is not a cosmetics, use only for the affected areas listed in the indications, and do not use on the face for the purpose of basic skin care, etc.

**INDICATIONS**

Dry skin accompanied by itching (xeroderma in the elderly/adults, dry skin in children)

**PRECAUTIONS FOR USE**

☒ The following precaution should be observed.

(If not observed, it may aggravate the current symptoms or increase the risk of adverse reactions.)

Do not apply to the following areas.

- (1) Around the eyes, mucous membranes (e.g. mouth, nasal cavity, vagina, etc.)
- (2) Inflammation or wounds
- (3) Severe skin sore or cracking of the skin

### Consultation

1. The following persons should consult a physician, pharmacist or registered salesperson before using the medicine.

- (1) Persons under treatment by a physician
- (2) Persons with a history of allergic symptoms (e.g. rash, redness, itching, contact dermatitis, etc.) or irritation symptoms due to any drug

2. The following symptoms may be adverse reactions to the medicine. If any of these symptoms occur, discontinue the use of the medicine immediately, and consult a physician, pharmacist or registered salesperson with this leaflet.

Affected body system	Symptoms
Skin	Contact dermatitis, rash, redness, itching, irritation (pain, feeling hot, tingling sensation), swelling, scab-like skin exfoliation

3. If symptoms do not improve after using the medicine for about two weeks, discontinue the use of the medicine, and consult a physician, pharmacist or registered salesperson with this leaflet.