


Brand name: Kansei																															
<p>RISK CATEGORIES Second-class OTC drug</p> <p>CHARACTERISTICS I This is a medicine to support liver function. Kansei is an herbal medicine originally formulated with 12 types of crude drugs including those to improve liver function, those to enhance bile secretion, and those with both effects. This medicine is recommended for persons who are diagnosed as having mild abnormal hepatic function at health examination, complete medical checkup, etc.</p>																															
	<p>ACTIVE INGREDIENTS 3 pouches of Kansei (6 g) contain 2 g of dried Kansei extract prepared from the following crude drugs.          Jujube Seed...1.46 g,          Ginseng...1.46 g          Gardenia Fruit...1.46 g,          Plantago Seed...1.46 g          Mulberry Bark...1.46 g,          Immature Orange...1.46 g          Amomum Seed...1.46 g,          Cinnamon Bark...0.37 g          Rhubarb...0.37 g,          Mume Fructus...1.46 g          Tachibana Pericarpium...1.46 g,          Artemisia Leaf...1.46 g          Inactive ingredients: Lactose, Carmellose Sodium</p> <p>&lt;Precautions&gt; As this medicine is manufactured from crude drugs, slight difference in color and taste may occur between each product, but will not affect the quality.</p>																														
<p>For further information:</p> <table border="1" style="width: 100%;"> <tr> <td>Distributed by:</td> <td colspan="3">TAIHO PHARMACEUTICAL CO., LTD.</td> </tr> <tr> <td>Manufactured and Distributed by:</td> <td colspan="3">Tatebayashi Shokakudo Co., Ltd.</td> </tr> <tr> <td>Dosage form</td> <td colspan="3">Granules</td> </tr> <tr> <td>Packaging unit</td> <td>2 g x 1 pouches</td> <td>2 g x 21 pouches</td> <td>2 g x 60 pouches</td> </tr> <tr> <td>Manufacturer's suggested retail price</td> <td>JPY 253 (JPY 230 excluding tax)</td> <td>JPY 3,163 (JPY 2,875 excluding tax)</td> <td>JPY 7,590 (JPY 6,900 excluding tax)</td> </tr> <tr> <td>JAN Code</td> <td>49684360</td> <td>4987117219104</td> <td>4987117219005</td> </tr> <tr> <td>Expiration</td> <td colspan="3">5 years</td> </tr> </table>				Distributed by:	TAIHO PHARMACEUTICAL CO., LTD.			Manufactured and Distributed by:	Tatebayashi Shokakudo Co., Ltd.			Dosage form	Granules			Packaging unit	2 g x 1 pouches	2 g x 21 pouches	2 g x 60 pouches	Manufacturer's suggested retail price	JPY 253 (JPY 230 excluding tax)	JPY 3,163 (JPY 2,875 excluding tax)	JPY 7,590 (JPY 6,900 excluding tax)	JAN Code	49684360	4987117219104	4987117219005	Expiration	5 years		
Distributed by:	TAIHO PHARMACEUTICAL CO., LTD.																														
Manufactured and Distributed by:	Tatebayashi Shokakudo Co., Ltd.																														
Dosage form	Granules																														
Packaging unit	2 g x 1 pouches	2 g x 21 pouches	2 g x 60 pouches																												
Manufacturer's suggested retail price	JPY 253 (JPY 230 excluding tax)	JPY 3,163 (JPY 2,875 excluding tax)	JPY 7,590 (JPY 6,900 excluding tax)																												
JAN Code	49684360	4987117219104	4987117219005																												
Expiration	5 years																														
<p>DOSAGE AND ADMINISTRATION Take the medicine with water or warm water between meals.</p> <table border="1" style="width: 100%;"> <thead> <tr> <th>Age</th> <th>Dose</th> <th>Daily dose</th> </tr> </thead> <tbody> <tr> <td>Adults (15 years and over)</td> <td>1 pouch (2 g)</td> <td>3 times</td> </tr> <tr> <td>Under 15 years</td> <td colspan="2">Do not use.</td> </tr> </tbody> </table> <p>Between meals mean 2 to 3 hours after meals.</p>				Age	Dose	Daily dose	Adults (15 years and over)	1 pouch (2 g)	3 times	Under 15 years	Do not use.																				
Age	Dose	Daily dose																													
Adults (15 years and over)	1 pouch (2 g)	3 times																													
Under 15 years	Do not use.																														
<p>INDICATIONS Abnormal hepatic function, hepatic hypertrophy, acute/chronic hepatitis, jaundice, cholecystitis</p>																															

## PRECAUTIONS FOR USE

☒ The following precaution should be observed.

(If not observed, it may aggravate the current symptoms or increase the risk of adverse reactions.)

Lactating mothers should avoid using the medicine or stop breastfeeding.

### Consultation

1. The following persons should consult a physician, pharmacist or registered salesperson before using this medicine.

- (1) Persons under treatment by a physician
- (2) Pregnant or possibly pregnant women
- (3) Persons who are physically weak (declining of strength or constitutional weakness)
- (4) Persons who have gastrointestinal weakness and are prone to diarrhea
- (5) Persons with a history of rash, redness, itching, etc. due to any drug
- (6) Persons who are taking the following drugs  
Cathartic drugs (laxatives)

2. The following symptoms may be adverse reactions to the medicine. If any of these symptoms occur, discontinue the use of the medicine immediately, and consult a physician, pharmacist or registered salesperson with this leaflet.

Affected body system	Symptoms
Skin	Rash, redness, itching
Gastrointestinal	Diarrhea accompanied by severe abdominal pain, abdominal pain

\* The following serious symptoms occur in rare cases. In such cases, consult a physician immediately.

Name of symptom	Symptoms
Mesenteric vein sclerosis	Abdominal pain, diarrhoea, constipation, abdominal distension, etc. occur repeatedly by long-term taking.

3. The following symptoms may be adverse reactions to the medicine. If any of these symptoms persist or get worse, discontinue the use of the medicine immediately, and consult a physician, pharmacist or registered salesperson with this leaflet.

Soft stool, diarrhea

4. If symptoms do not improve after using the medicine for about one month, discontinue the use of this medicine, and consult a physician, pharmacist or registered salesperson with this leaflet.
5. When the medicine is to be used continuously for a long term, consult a physician, pharmacist or registered salesperson.

## CONTACT

For inquiries, please contact the shop where you purchased the product or the following:

Distributor Contact: Customer Relations Office, Taiho Pharmaceutical Co., Ltd.

1-27 Kandnishiki-cho, Chiyoda-ku, Tokyo, 101-8444, Japan

Phone: 0120-4527-66 Business hours: 9:00-17:00 (Excluding weekends and holidays)

Website: <https://www.taiho.co.jp/>

Manufactured and Distributed by : Tatebayashi Shokakudo Co., Ltd.