

Brand name: Solmack 5

**RISK CATEGORIES**  
Designated quasi-drug

**CHARACTERISTICS**

I Gastrointestinal medicine containing crude drugs  
The first product of Solmack series to be taken before meals  
Easy-to-take refreshing plum flavor  
It contains a maximum of 2,000 mg of turmeric (maximum amount within the standard range for registered quasi-drug: equivalent to the amount of crude drug per 50 mL bottle)  
The medicine contains perilla herb and red ginseng which help improve gastrointestinal function.  
A designated quasi-drug "Solmack 5" contains five types of crude drugs.  
It exerts an excellent effect on stomach discomfort and feeling queasy due to overeating or overdrinking.



**ACTIVE INGREDIENTS**

Per bottle (50 mL)  
Turmeric Extract 200 mg (equivalent to 2,000 mg of turmeric)  
Red Ginseng Extract 83.3 mg (equivalent to 333.2 mg of red ginseng)  
Perilla Herb Extract 37 mg (equivalent to 333 mg of perilla herb)  
Clove Oil 5 mg  
Dipotassium Glycyrrhizinate 7.2 mg

Inactive ingredients: Glycerin, Fructose, Propylene Glycol, Glucose, Povidone, Citric Acid Hydrate, Polyoxyethylene Hydrogenated Caster Oil, l-Menthol, Sodium Benzoate, pH Adjuster, Vanillin, Flavor

● Shake well before use. As this medicine contains crud drugs, it may cause precipitation.

**INDICATIONS**

Stomach discomfort or feeling queasy (sickness, churning of stomach, hangover, drunken sickness, queasy, nausea) due to overeating or overdrinking

For further information:

Manufactured and Distributed by:	TAIHO PHARMACEUTICAL CO., LTD.	
Dosage form	Liquid	
Packaging unit	50 mL x 1 bottle	50 mL x 8 bottles
Manufacturer's suggested retail price	JPY 385 (JPY 350 excluding tax)	JPY 3,080 (JPY 2,800 excluding tax)
JAN Code	45144653	4987117396065
Expiration	3 years	

**DOSAGE AND ADMINISTRATION**

Adults (15 years and over): Take 1 bottle (50 mL) before meals once daily.

- Do not use in children (under 15 years).
- Follow the dosing instruction.

**PRECAUTIONS FOR USE**

■ Consultation

1. In the following cases, discontinue the use of the medicine immediately, and consult a physician or pharmacist with this medicine.

- (1) If rash occurs after using the medicine.
- (2) If symptoms do not improve after using the medicine for a while.

2. Diarrhea may occur. If the symptom persists or gets worse, discontinue the use of the medicine, and consult a physician or pharmacist.

## CONTACT

For inquiries, please contact the shop where you purchased the product or the following:

Manufacturer and distributor Contact:

Customer Relations Office, Taiho Pharmaceutical Co., Ltd. 1-27 Kandanishiki-cho, Chiyoda-ku, Tokyo, 101-8444,  
Japan

Phone: 0120-4527-66 Business hours: 9:00-17:00 (Excluding weekends and holidays)

Website: <https://www.taiho.co.jp/>