

Brand name: TIOVITA Drink 2000

RISK CATEGORIES
Designated quasi-drug

CHARACTERISTICS
I Vitamin health drink
- “Effective” and “Good Taste” Green Tiovita! with a catchphrase “One Love to Share”!
- It contains 2,000 mg of taurine as an active ingredient (twice more than Tiovita Drink).
- Easy-to-take “cassis flavor”
- It only has “72 kcal”.



ACTIVE INGREDIENTS
Per 100 mL bottle
Taurine 2,000 mg, Inositol 50 mg, Nicotinamide 20 mg, Thiamine Nitrate (Vitamin B₁) 5 mg, Riboflavin Sodium Phosphate (Vitamin B₂) 5 mg, Pyridoxine Hydrochloride (Vitamin B₆) 5 mg, Carnitine Hydrochloride 100 mg, Anhydrous Caffeine 50 mg

Inactive ingredients: White Soft Sugar, Fructose, Glucose, DL-Malic Acid, Propyl Gallate, Sodium Benzoate, pH Adjuster, Vanillin, Flavor
● Urine may turn yellow because of vitamin B₂ contained in the medicine. It is nothing to worry about.

INDICATIONS
○ Nutritional fortification
○ Weak constitution
○ Supply of nutrients in the following cases: physical fatigue, during and after illness, loss of appetite, nutritional disorders, febrile debilitating illness, during pregnancy or lactation

For further information:

Manufactured and Distributed by:	TAIHO PHARMACEUTICAL CO., LTD.			
Dosage form	Liquid			
Packaging unit	100 mL x 1 bottle	100 mL x 3 bottles	100 mL x 10 bottles	100 mL x 50 bottles
Manufacturer's suggested retail price	JPY 228 (JPY 207 excluding tax)	JPY 683 (JPY 621 excluding tax)	JPY 2,277 (JPY 2,070 excluding tax)	JPY 11,385 (JPY 10,350 excluding tax)
JAN Code	4987117439106	4987117439205	4987117439113	4987117439120
Expiration	3 years			

DOSAGE AND ADMINISTRATION
Adults (15 years and over): Take 1 bottle (100 mL) once daily.
● Do not use in children (under 15 years).
● Follow the dosing instruction. (Please pay attention to overdose if other products containing vitamins are concomitantly used.)

PRECAUTIONS FOR USE
■ Consultation
1. In the following cases, discontinue the use of the medicine immediately, and consult a physician or pharmacist with this medicine.
(1) If rash occurs after using the medicine.
(2) If symptoms do not improve after using the medicine for a while.
2. Diarrhea may occur. If the symptom persists or gets worse, discontinue the use of the medicine, and consult a physician or pharmacist.

CONTACT

For inquiries, please contact the shop where you purchased the product or the following:

Manufacturer and distributor Contact:

Customer Relations Office, Taiho Pharmaceutical Co., Ltd.

1-27 Kandanshiki-cho, Chiyoda-ku, Tokyo, 101-8444, Japan

Phone: 0120-4527-66 Business hours: 9:00-17:00 (Excluding weekends and holidays)

Website: <https://www.taiho.co.jp/>