

Brand name: TIOVITA GOLD 2000

**RISK CATEGORIES**

Second-class OTC drug

**CHARACTERISTICS**

I Nutrient tonic drink

- Taurine 2,000 mg and crude drug extracts exert effects!
- The medicine contains royal jelly and epimedium herb.(For further information, please see ACTIVE INGREDIENTS.)
- "Apricot flavor" which is easy to take for women
- It only has "52 kcal".



**ACTIVE INGREDIENTS**

Per 50 mL bottle

Taurine 2,000 mg, Royal Jelly Extract 0.15 mL (equivalent to 100 mg of royal jelly), Epimedium Herb Fluidextract 0.1 mL (equivalent to 100 mg of epimedium herb), Thiamine Nitrate (Vitamin B<sub>1</sub>) 5 mg, Riboflavin Sodium Phosphate (Vitamin B<sub>2</sub>) 5 mg, Pyridoxine Hydrochloride (Vitamin B<sub>6</sub>) 5 mg, Nicotinamide 30 mg, Anhydrous Caffeine 50 mg (≤ 0.47 mL of alcohol)

Inactive ingredients: Fructose, Citric Acid, Propyl Gallate, Ginseng Fluidextract, Alcohol, Sodium Benzoate, Paraben, pH Adjuster, Vanillin, Flavor

- (1) Shake well before use. As this medicine contains crudrugs, it may cause precipitation.
- (2) Urine may turn yellow because of vitamin B<sub>2</sub> contained in the medicine. It is nothing to worry about.

**INDICATIONS**

Nutritional fortification, weak constitution, and supply of nutrients in the following cases: physical fatigue, during and after illness, loss of appetite, nutritional disorders, febrile debilitating illness, during pregnancy or lactation

For further information:

Manufactured and Distributed by:	TAIHO PHARMACEUTICAL CO., LTD.	
Dosage form	Liquid	
Packaging unit	50 mL x 1 bottle	50 mL x 10 bottles
Manufacturer's suggested retail price	JPY 418 (JPY 380 excluding tax)	JPY 4,180 (JPY 3,800 excluding tax)
JAN Code	4987117437409	4987117437416
Expiration	3 years	

**DOSAGE AND ADMINISTRATION**

Adults (15 years and over): Take 1 bottle (50 mL) once daily.

- Do not use in children (under 15 years).
- Follow the dosing instruction.

**PRECAUTIONS FOR USE**

■ Consultation

1. Diarrhea may occur after using the medicine. If the symptom persists or gets worse, discontinue the use of the medicine, and consult a physician, pharmacist or registered salesperson with this medicine.
2. If symptoms do not improve after using the medicine for a while, discontinue the use of the medicine, and consult a physician, pharmacist or registered salesperson with this medicine.

**CONTACT**

For inquiries, please contact the shop where you purchased the product or the following:

Manufacturer and distributor Contact:

Customer Relations Office, Taiho Pharmaceutical Co., Ltd.

1-27 Kandanshiki-cho, Chiyoda-ku, Tokyo, 101-8444, Japan

Phone: 0120-4527-66 Business hours: 9:00-17:00 (Excluding weekends and holidays)

Website: <https://www.taiho.co.jp/>